

References

- (1) Thomson, J. J. *Phil. Mag.* **1911**, 21, 225.
- (2) Aston, F. W. Neon. *Nature* **1919**, 104, 334.
- (3) Aston, F. W. A positive-ray spectrograph. *Phil. Mag.* **1919**, 38, 707-715.
- (4) Aston, F. W. A new mass spectrograph and the whole-number rule (Bakerian Lecture). *Proc. Roy. Soc. (London)* **1927**, A115, 487-514.
- (5) Aston, F. W. A second-order focusing mass spectrograph and isotopic weights by the doublet method. *Proc. Roy. Soc. (London)* **1937**, A163, 391-404.
- (6) Dempster, A. J. A New Method of Positive Ray Analysis. *Phys. Rev.* **1918**, 11, 316.
- (7) Dempster, A. G. Positive-Ray Analysis of Potassium, Calcium and Zinc. *Phys. Rev.* **1922**, 20, 631-638.
- (8) Dempster, A. J. New methods in mass spectroscopy. *Proc. Am. Phil. Soc.* **1935**, 75, 755-767.
- (9) Bainbridge, K. T.; Jordan, E. B. Mass-spectrum analysis. 1. The mass spectrograph. 2. The existence of isobars of adjacent elements. *Physical Review* **1936**, 50, 282-296.
- (10) Mattauch, J.; Herzog, R. Mass spectrograph. *Zeitschrift fuer Physik* **1934**, 89, 786-795.
- (11) Mattauch, J. A double-focusing mass spectrograph and the masses of N15 and O18. *Physical Review* **1936**, 50, 617-623.
- (12) Hintenberger, H.; Konig, L. A. Mass spectrometer with complete double focusing in second order. *Zeitschrift fuer Naturforschung* **1957**, 12a, 773-785.
- (13) Okuda, T.; Ogata, K.; Aoki, K.; Sugawara, Y. The isotopic weights of chlorine, argon and iron by the doublet method. *Physical Review* **1940**, 58, 578-579.
- (14) Ogata, K.; Matsuda, H. The mass differences C12H14-O16, C12H12-N14, and C12H13N15. *Physical Review* **1951**, 83, 180-181.
- (15) Matsuda, H. Focusing properties of a large mass spectrograph newly constructed at Osaka University. *Journal of the Physical Society of Japan* **1956**, 11, 183-191.
- (16) Johnson EG; Nier AO Angular aberrations in sector-shaped electromagnetic lenses for focusing beams of charged particles. *Phys.Rev.* **1953**, 91, 10-17.
- (17) Nier, A. O.; Roberts, T. R. The determination of atomic mass doublets by means of a mass spectrometer. *Phys. Rev.* **1951**, 81, 507.